arXiv:1212.0357v1 [hep-ph] 3 Dec 2012

Ballpark prediction for the hadronic light-by-light contribution to the muon $(g-2)_\mu$

Pere Masjuan* and Marc Vanderhaeghen
Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

(Dated: December 4, 2012)

Using the momentum dependence of the dressed quark mass and the well-known formulae for the mass dependent quark loop contribution to the light-by-light scattering insertions, we compute the hadronic light-by-light contribution to the the muon anomalous magnetic moment. We ascribe for the first time a systematic error on the calculation.

PACS numbers: 13.40.Em,14.60.Ef

The anomalous magnetic moment of the muon is one of the most accurately measured quantities in particle physics. Any deviation from its prediction in the Standard Model of particle physics is a very promising signal of new physics.

The present world average experimental value of its deviation from the Dirac value, i.e., $a_\mu = (g_\mu - 2)/2$, is given by $a_\mu^{EXP} = 11659208.9(6.3) \times 10^{-10}$ [1, 2]. This impressive result is still limited by statistical errors, and a proposal to measure the muon $(g-2)_\mu$ to a precision of 1.6×10^{-10} has recently been submitted to FNAL [3].

At the level of the experimental accuracy, the QED contributions to a_μ from photons and leptons alone are very well known. Recently the calculation has been completed up to the fifth order $\mathcal{O}(\alpha_{em}^5)$ in the fine-structure constant α_{em}, giving the result $11658471.885(4) \times 10^{-10}$ [4].

The main uncertainties at present in the Standard Model calculation for $(g-2)_\mu$ originate from the hadronic vacuum polarization (HVP) as well as hadronic light-by-light scattering (HLBL) corrections. We show the present estimates and their uncertainties for the QED, HVP, HLBL, and the electroweak (EW) corrections in Table I.

TABLE I. Standard Model contributions to $(g-2)_\mu$.

Contribution	Result in 10^{-10} units	Ref.
QED(leptons)	11658471.885 ± 0.004	[4]
HVP(leading order)	692.3 ± 4.2	[5]
HVP(higher order)	-9.84 ± 0.07	[6]
HLBL	11.6 ± 4.0	[7]
EW	15.4 ± 0.2	[8]
Total	11659181.3 ± 5.8	

The existing discrepancy between the experimental value for $(g-2)_\mu$ and its Standard Model prediction stands at about 3σ.

In order to interpret the upcoming new experiment at FNAL, with an anticipated precision of 1.6×10^{-10}, there is an urgent need to improve on both the HVP as well as the HLBL contributions. The accuracy of the HVP contribution depends on the statistical error of the experimental data for the e^+e^- annihilation cross-section into hadrons. With future experiments, in particular at BES-III [9], one foresees this error to quantitatively decrease. The HLBL cannot be directly related to any measurable cross section however, and requires the knowledge of Quantum Chromodynamics (QCD) contributions at all energy scales. Since this is not known yet, one needs to rely on hadronic models to compute it. Such models introduce some systematic errors which are difficult to quantify.

The main motivation of this work is to provide a ballpark prediction with a judicious error estimate for the HLBL scattering based on a duality argument between the hadronic degrees of freedom and the well-known quark loop contribution.

Such a duality estimate can be obtained by invoking a particular regime of QCD where one knows how to perform the quark loop integral responsible for the a_μ^{HLBL} (Fig. 1). This is the large-N_c of QCD [10, 11] where a quark-hadron duality is accounted for considering that hadronic amplitudes are described by an infinite set of non-interacting and non-decaying resonances. As shown in Ref. [12, 13], the large-N_c limit provides a very useful framework to approach this problem.

Using the large-N_c counting and also the chiral counting, it was proposed in [12] to split the diagram of Fig. 1 into a set of different contributions where the numerically dominant contribution arises from the pseudo-scalar exchange diagram shown in Fig. 2 [7].

The large-N_c approach however has two shortcomings: firstly, the assumption of pion-exchange dominance implies that the remaining pieces are small enough to justify their omission. Although this seems reasonable [13], it might lead to an underestimation of the error. Secondly, calculations carried out in the large-N_c limit demand an infinite set of resonances for computing any quantity. As such sum is not known in practice, one ends up truncating the spectral function in a resonance saturation scheme, the so-called Minimal Hadronic Approximation [14]. The resonance masses used in each calculation are then taken as the physical ones from PDG [15] instead of the corresponding masses in the large-N_c limit. Both problems might lead to large systematic errors not included so far

---

* Corresponding author: masjuan@kph.uni-mainz.de



FIG. 1. Quark loop diagram with running quark mass.



FIG. 2. Pion-pole contribution to HLBL scattering. The shaded blobs represent the Form Factor $F_{\pi^0\gamma^*\gamma^*}(q_1^2, q_2^2)$.

[16, 17].

It was pointed out in Ref. [16] that, in the large-N_c framework, the Minimal Hadronic Approximation can be understood from the mathematical theory of Pade Approximants (PA) to meromorphic functions. Obeying the rules from this mathematical framework, one can compute the desired quantities in a model-independent way and even be able to ascribe a systematic error to the approach [18]. One interesting detail from this theory [19] is that given a low-energy expansion of a meromorphic function, a PA sequence converges much faster than a rational function with the poles fixed in advance (such as the common hadronic models used so far for evaluating the HLBL), especially when the correct large Q^2 behavior is imposed.

To perform a quark-hadron duality estimate for the HLBL contribution to a_μ, we now discuss the direct contribution from the quark-loop diagram. The diagram with a light-quark running, with a mass of a few MeV, is only valid in the regime where quarks are not confined, so in the perturbative QCD regime. This is not the region dominating the loop integral since we know that the momentum circulating the loop covers the low- and high-energy ranges. From quark models, one can use a constituent quark mass model with mass around 300 MeV as a rough estimate (see [20] for a recent computation). This however, fails to reproduce the perturbative QCD result at high energies. One, then, desires a running momentum dependent mass to perform such integration.



FIG. 3. Momentum dependent dressed quark mass, $M(Q)$, obtained as a solution of the gap equation within the Dyson-Schwinger formalism. The horizontal line represents the muon mass. The band corresponds to an error of 10% on top of the chiral limit (solid red line).

Ideally, lattice QCD calculations could be able to calculate such momentum dependent quark masses. As at present lattice QCD calculations are not yet fully feasible for physical pion masses, our proposal is to use the dressed-quark mass function $M(Q)$ in the chiral limit computed within the Dyson-Schwinger equation (DSE) framework. As shown in Fig. 3, the solution of the gap equation for a renormalized dressed-quark propagator using DSE provides us with the desired momentum dependent quark-mass function [21, 22]. We add a 10% relative error as an estimate for the extrapolation from the chiral limit to the physical light quark mass regime [23].

To determine the average momentum of the quark running in the loop on Fig. 1, we employ for our duality argument a hadronic model. Such model yields the average momenta entering the momentum dependent dressed mass $M(Q_1)$ and $M(Q_2)$, where $Q_{1,2}$ refer to the momentum labels as indicated in Fig. 2. With this mass, one can use the well-known formulae for the a_μ^{HLBL} for spin$-1/2$ fermions computed in [24–26]:

$$a_\mu^{HLBL}(M(Q)) = \left(\frac{\alpha}{\pi}\right)^3 N_c \left(\sum_{q=u,d,s} Q_q^4\right)\left[\left(\frac{3}{2}\zeta(3) - \frac{19}{16}\right)\frac{m_\mu^2}{M(Q)^2} + \mathcal{O}\left(\frac{m_\mu^4}{M(Q)^4}\log^2\frac{m_\mu^2}{M(Q)^2}\right)\right]. \quad (1)$$

In Eq. (1) we display the result up to first order in $(m_\mu^2/M(Q)^2)$ but we include all terms up to the fourth order in our numerical calculations [7].

For our evaluation we will project the hadronic content of the quark loop of Fig. 1 onto the dominant hadronic piece of the HLBL. This is the pion-exchange contribution considered in [27]. We will use, instead of a hadronic model for the transition Form Factors (FF), a sequence of rational approximants [16] build up from the low-energy expansion of the pion FF obtained in [17] after a fit to the experimental data [28–31], to minimize a model dependence.

In practice that means to use for the FFs (blobs in Fig. 2) the expression:

$$F^{P01}_{\pi\gamma^*\gamma^*}(q_1^2, q_2^2) \;=\; P_1^0(q_1^2, q_2^2) \;= a\frac{b}{q_1^2 - b}\frac{b}{q_2^2 - b}\,, \qquad (2)$$

where the free parameters are matched at low energies with the results in [17]: a is fixed by $\Gamma_{\pi\to\gamma\gamma} = 7.82(24)$eV from the PrimEx Collaboration [32]; and b by a matching to the slope $a_\pi = 0.0324(22)$ [17]. The results for the average momenta running in the quark loop in Fig. 1 using the FFs of Eq. (2) are shown in Table II where in parenthesis we quote the errors from the input uncertainties.

The convergence of the PA sequence to a meromorphic function is guaranteed by Pomerenke theorem [33]. The problem is to know how fast this convergence is and also how to ascribe a systematic error on each element of that sequence. For the particular case of a meromorphic function (such as a Green's function in large-N_c QCD), the simplest way of evaluating a systematic error is by comparing the difference between two consecutive elements on the PA sequence [18].

In our approach to the FF, we evaluate the systematic error by computing a second element on the PA sequence [18] and compare it with the result using Eq. (2). The second element is:

$$F^{P12}_{\pi\gamma^*\gamma^*}(q_1^2, q_2^2) \;= \\ = P_2^1(q_1^2, q_2^2) \;= \frac{a + bq_1^2}{(q_1^2 - c)(q_1^2 - d)}\frac{a + bq_2^2}{(q_2^2 - c)(q_2^2 - d)}\,, \qquad (3)$$

with four coefficients to be matched with $\Gamma_{\pi\to\gamma\gamma}$, the slope a_π, the curvature of the pion FF $b_\pi = 1.06(26) \times 10^{-3}$ [18] and the first vector mesonresonance $M_\rho = 0.7755(3)$ GeV [15]. The results for the average momenta running in the quark loop in Fig. 1 using the approximant of Eq. (3) are shown in Table II where again the errors are due to the input ones.

The similarity of the results obtained within both approximants indicates that the low-energy region (up to $1 - 2$ GeV) dominates the contribution to a_μ^{HLBL}. To evaluate the error on our approximation we look for the maximum of the difference in the region up to 1 GeV between the $P_1^0(q_1^2, q_2^2)$ and $P_2^1(q_1^2, q_2^2)$ as explained in Ref. [18]. Of course, this difference depends on the energy, and grows as the energy increases. At 1 GeV, the relative difference is about 5%, and we take this error as a conservative estimate of the error on the whole low-energy region. We should add this error to the a_μ^{HLBL} results shown in Table II.

TABLE II. Collected results for the average momenta running in the quark loop in Fig. 1, its corresponding mass $M(Q)$ in Fig. 3, and the a_μ^{HLBL} result in accordance to Eq. (1), Fig. 4, for both $P_1^0(Q_1^2, Q_2^2)$ and $P_2^1(Q_1^2, Q_2^2)$.

		$\langle Q_i \rangle$	$M(Q_i)$GeV	$a_\mu^{HLBL} \times 10^{10}$
$P_1^0(Q_1^2, Q_2^2)$	Q_1	0.42(1)GeV	0.186(3)	12.5(3)
	Q_2	0.29(1)GeV	0.224(2)	9.0(1)
$P_2^1(Q_1^2, Q_2^2)$	Q_1	0.43(2)GeV	0.184(6)	12.7(7)
	Q_2	0.30(1)GeV	0.222(3)	9.2(2)



FIG. 4. a_μ^{HLBL} results using the expansion in Eq. (1) at order $(m_\mu^2/M(Q)^2)^6$ (dashed blue) and at order $(m_\mu^2/M(Q)^2)^8$ (solid black) in terms of the running quark mass $M(Q)$. The red point indicates the exact result for the point $M(Q) = m_\mu$. The horizontal band shows the ballpark result from Eq. (4).

To quote a final number for a_μ^{HLBL} implies to consider several sources of error:

- Firstly, we have the error coming from the experimental inputs used to build up our approximants, which arises mainly from the fit to the experimental data on the FF. With the new forthcoming experimental data on that FF at BES-III [9] we intuit lower input errors on our results.

- Secondly, we have a 10% error due to the departure from the chiral limit shown in Fig. 3. This error is absolutely asymmetric and moves the $M(Q)$ values in Table II upwards. That implies a decrease of about 15% on a_μ^{HLBL} (down to 10.6 and 7.6 in units of 10^{-10} for Q_1 and Q_2 respectively). We are using the DSE to obtain the momentum dependent dressed quark masses. Once the lattice calculations will reach the physical quark mass values, this 10% error will disappear.

- Furthermore, we also have a systematic error from the PA sequence used. We estimate this to be around 5%. That implies an error on a_μ^{HLBL} about ± 0.6 for Q_1 and ± 0.5 for Q_2 in units 10^{-10}. Since both $P_1^0(q_1^2, q_2^2)$ and $P_2^1(q_1^2, q_2^2)$ give almost the same results for a_μ^{HLBL}, no extra error due to the difference between them should be included on a_μ^{HLBL}. We should remark that the FFs employed here, although constrained by the experimental data, do not have the correct behavior when both photon virtualities (q_1 and q_2) are very large where a behavior $\sim 1/(q_1^2 + q_2^2)$ is predicted [34–36]. This fact does not affect our calculation since a_μ^{HLBL} is very largely dominated by the low-energy region.

- The last source of error comes from the evaluation of a_μ^{HLBL} using the order $(m_\mu^2/M(Q)^2)^8$ in Eq. (1) instead of the full result. The difference is so smooth, Fig. 4, that no extra error should be considered so far.

Our ballpark estimation lies then in the range:

$$a_\mu^{HLBL} = [7.6(4) \div 12.5(7)] \times 10^{-10} \,, \qquad (4)$$

where the error in parenthesis is the combined systematic and input errors as commented above and the two numbers represent the range due to the two momenta considered in our computations.

In summary, we presented a ballpark estimate of the hadronic light-by-light scattering contribution to the $(g-2)_\mu$ based on a duality argument, and estimated the average momentum flowing in the quark loop diagram of Fig. 1 from a hadronic model. This average momentum then allowed us to calculate the momentum dependent quark mass in the quark-loop result from a_μ^{HLBL}. Most of the recent phenomenological calculations of the a_μ^{HLBL} fall into the range obtained in this work ([13, 20, 27, 37–39]).

We would like to thank StanBrodsky for helpful discussions and to I. Cloët for providing us with the momentum dependent dressed quark mass function shown in Fig. 3. The present work is supported by the Deutsche Forschungsgemeinschaft DFG through the Collaborative Research Center "The Low-Energy Frontier of the Standard Model" (SFB 1044).

[1] G. Bennett *et al.* (Muon g-2 Collaboration), Phys.Rev.Lett. **92**, 161802 (2004), arXiv:hep-ex/0401008 [hep-ex].

[2] G. Bennett *et al.* (Muon G-2 Collaboration), Phys.Rev. **D73**, 072003 (2006), arXiv:hep-ex/0602035 [hep-ex].

[3] R. Carey, K. Lynch, J. Miller, B. Roberts, W. Morse, *et al.*, (2009).

[4] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys.Rev.Lett. **109**, 111808 (2012), arXiv:1205.5370 [hep-ph].

[5] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, Eur.Phys.J. **C71**, 1515 (2011), arXiv:1010.4180 [hep-ph].

[6] K. Hagiwara, R. Liao, A. D. Martin, D. Nomura, and T. Teubner, J.Phys. **G38**, 085003 (2011).

[7] F. Jegerlehner and A. Nyffeler, Phys.Rept. **477**, 1 (2009), arXiv:0902.3360 [hep-ph].

[8] A. Czarnecki, W. J. Marciano, and A. Vainshtein, Phys.Rev. **D67**, 073006 (2003), arXiv:hep-ph/0212229 [hep-ph].

[9] D. Asner, T. Barnes, J. Bian, I. Bigi, N. Brambilla, *et al.*, Int.J.Mod.Phys. **A24**, S1 (2009), arXiv:0809.1869 [hep-ex].

[10] G. 't Hooft, Nucl. Phys. **B72**, 461 (1974).

[11] E. Witten, Nucl. Phys. **B160**, 57 (1979).

[12] E. de Rafael, Phys.Lett. **B322**, 239 (1994), arXiv:hep-ph/9311316 [hep-ph].

[13] J. Prades, E. de Rafael, and A. Vainshtein, (2009), arXiv:0901.0306 [hep-ph].

[14] S. Peris, M. Perrottet, and E. de Rafael, JHEP **9805**, 011 (1998), arXiv:hep-ph/9805442 [hep-ph].

[15] J. Beringer *et al.* (Particle Data Group), Phys.Rev. **D86**, 010001 (2012).

[16] P. Masjuan and S. Peris, JHEP **0705**, 040 (2007), arXiv:0704.1247 [hep-ph].

[17] P. Masjuan, Phys.Rev. **D86**, 094021 (2012), arXiv:1206.2549 [hep-ph].

[18] P. Masjuan and S. Peris, Phys.Lett. **B686**, 307 (2010), arXiv:0903.0294 [hep-ph].

[19] P. Masjuan Queralt, Ph.D. thesis, Universitat Autònoma de Barcelona (2010), arXiv:1005.5683 [hep-ph].

[20] D. Greynat and E. de Rafael, JHEP **1207**, 020 (2012), arXiv:1204.3029 [hep-ph].

[21] M. Bhagwat, M. Pichowsky, C. Roberts, and P. Tandy, Phys. Rev. **C68**, 015203 (2003).

[22] M. Bhagwat, I. Cloet, and C. Roberts, (2007), arXiv:0710.2059 [nucl-th].

[23] P. O. Bowman, U. M. Heller, D. B. Leinweber, and A. G. Williams, Nucl.Phys.Proc.Suppl. **119**, 323 (2003), arXiv:hep-lat/0209129 [hep-lat].

[24] J. Aldins, T. Kinoshita, S. J. Brodsky, and A. Dufner, Phys.Rev.Lett. **23**, 441 (1969).

[25] J. Aldins, T. Kinoshita, S. J. Brodsky, and A. Dufner,

Phys.Rev. **D1**, 2378 (1970).

[26] S. Laporta and E. Remiddi, Phys.Lett. **B301**, 440 (1993).

[27] M. Knecht and A. Nyffeler, Phys.Rev. **D65**, 073034 (2002), arXiv:hep-ph/0111058 [hep-ph].

[28] H. J. Behrend *et al.* (CELLO), Z. Phys. **C49**, 401 (1991).

[29] J. Gronberg *et al.* (CLEO), Phys. Rev. **D57**, 33 (1998), hep-ex/9707031.

[30] B. Aubert *et al.* (BABAR Collaboration), Phys.Rev. **D80**, 052002 (2009), arXiv:0905.4778 [hep-ex].

[31] S. Uehara *et al.* (Belle Collaboration), Phys.Rev. **D86**, 092007 (2012), arXiv:1205.3249 [hep-ex].

[32] I. Larin *et al.* (PrimEx Collaboration), Phys.Rev.Lett. **106**, 162303 (2011), arXiv:1009.1681 [nucl-ex].

[33] C. Pommerenke, J. Math. Anal. Appl. **41**, 775 (1973).

[34] F. del Aguila and M. Chase, Nucl.Phys. **B193**, 517 (1981).

[35] E. Braaten, Phys.Rev. **D28**, 524 (1983).

[36] S. J. Brodsky and G. P. Lepage, Adv.Ser.Direct.High Energy Phys. **5**, 93 (1989).

[37] M. Hayakawa and T. Kinoshita, Phys.Rev. **D57**, 465 (1998), arXiv:hep-ph/9708227 [hep-ph].

[38] J. Bijnens, E. Pallante, and J. Prades, Nucl.Phys. **B626**, 410 (2002), arXiv:hep-ph/0112255 [hep-ph].

[39] K. Melnikov and A. Vainshtein, Phys.Rev. **D70**, 113006 (2004), arXiv:hep-ph/0312226 [hep-ph].